



## CANCER BURDEN

### In 2000, the American Cancer Society estimates<sup>1</sup>

- 1,220,100 new cancer cases will be diagnosed in the United States, including 27,900 in Indiana.
- 552,200 cancer deaths will occur in the United States, including 12,600 in Indiana.

The average annual age-adjusted mortality rate for cancer deaths<sup>2</sup> per 100,000 persons

**Indiana: 177.6 National: 170.1**

*Indiana ranks 19<sup>th</sup> highest overall in cancer mortality rates among the 50 states and Washington, D.C.*

### LUNG CANCER

#### In 2000, the American Cancer Society estimates<sup>1</sup>

- 4,000 new cases of lung cancer will be diagnosed among men and women in Indiana.
- 3,900 men and women will die of lung cancer in Indiana.

The average annual age-adjusted mortality rates for lung cancer deaths<sup>2,3</sup> per 100,000 persons, by race, 1992–1996

	Indiana	National
Overall	55.3	49.5
White	54.6	49.2
Black	70.5	60.0
Hispanic	18.6	19.5
Asian/Pacific Islander	-	23.9
American Indian/Alaska Native	-	29.5

### BREAST CANCER

#### In 2000, the American Cancer Society estimates<sup>1</sup>

- 4,200 new cases of breast cancer will be diagnosed among women in Indiana.
- 900 women will die of breast cancer in Indiana.

The average annual age-adjusted mortality rates for breast cancer deaths<sup>2,3</sup> per 100,000 persons, by race, 1992–1996

	Indiana	National
Overall	25.8	25.4
White	25.5	25.1
Black	31.9	31.3
Hispanic	-	14.9
Asian/Pacific Islander	-	11.3
American Indian/Alaska Native	-	12.8

### COLORECTAL CANCER

#### In 2000, the American Cancer Society estimates<sup>1</sup>

- 3,100 new cases of colorectal cancer will be diagnosed among men and women in Indiana.
- 1,300 men and women will die of colorectal cancer in Indiana.

The average annual age-adjusted mortality rates for colorectal cancer deaths<sup>2,3</sup> per 100,000 persons, by race, 1992–1996

	Indiana	National
Overall	18.8	17.5
White	18.3	17.1
Black	27.6	22.9
Hispanic	8.8	10.3
Asian/Pacific Islander	-	10.9
American Indian/Alaska Native	-	9.9

### PROSTATE CANCER

#### In 2000, the American Cancer Society estimates<sup>1</sup>

- 3,900 new cases of prostate cancer will be diagnosed among men in Indiana.
- 700 men will die of prostate cancer in Indiana.

The average annual age-adjusted mortality rates for prostate cancer deaths<sup>2,3</sup> per 100,000 men, by race, 1992–1996

	Indiana	National
Overall	25.4	25.6
White	23.9	23.4
Black	52.5	54.8
Hispanic	-	16.6
Asian/Pacific Islander	-	10.8
American Indian/Alaska Native	-	14.7

<sup>1</sup> Source: American Cancer Society Facts and Figures, 2000. Estimates exclude about 1.3 million cases of basal and squamous cell skin cancers and in situ cancers, except urinary bladder, that will be diagnosed in 2000. Lung cancer rates include bronchus cancer. State death totals between 51-99 were rounded to 100.

<sup>2</sup> Source: CDC National Center for Health Statistics, vital statistics data, underlying cause of death, 1992–1996.

<sup>3</sup> Some data were excluded because there were 75,000 or fewer persons in the denominator or 20 or fewer deaths in the numerator.